

Overview of Stroud Water Research Center and Current Research on PFAS

Citizenship Advisory Council Meeting
PA Department of Environmental Protection




White Clay Creek, PA



Assistant Research Scientist

My research focuses on understanding transport and transformation of chemical substances at the watershed scale





- 1. Stroud Water Research Center
- 2. Current research on PFAS
- 3. Questions and Discussion





We advance knowledge and stewardship of freshwater systems through global research, education, and watershed restoration.






Guided by Science

The Stroud Center's independence, nonprofit status, and non-advocacy position ensure work is guided by science, not interests or agendas.

This fosters trust, credibility, and collaboration, empowering diverse partners and decision-makers with unbiased insights and allowing for nimble, mission-driven action focused on advancing knowledge and stewardship of fresh water.



- Independent 501(c)(3) nonprofit organization
- Located in Avondale, Pennsylvania






- Independent 501(c)(3) nonprofit organization
- Located in Avondale, Pennsylvania
- Founded in 1967 as a field station of the Natural Academy of Sciences of Philadelphia





- Independent 501(c)(3) nonprofit organization
- Located in Avondale, Pennsylvania
- Founded in 1967 as a field station of the Natural Academy of Sciences of Philadelphia
- On the banks of the East Branch of White Clay Creek



White Clay Creek Watershed, tributary of the Delaware



What We Do

Freshwater Research



Environmental Education




Watershed Restoration






Watershed-scale approach





Watershed-scale approach





Research Groups






- White Clay Creek is our natural laboratory
 - National Science Foundation designations:
 - 1981: Experimental Ecological Reserve
 - 1998: Long-term research in environmental biology site
 - River research at many spatial and temporal scales





- White Clay Creek is our natural laboratory
 - National Science Foundation designations:
 - 1981: Experimental Ecological Reserve
 - 1998: Long-term research in environmental biology site
 - River research at many spatial and temporal scales
 - Decades: forest buffer restoration





- White Clay Creek is our natural laboratory
 - National Science Foundation designations:
 - 1981: Experimental Ecological Reserve
 - 1998: Long-term research in environmental biology site
 - River research at many spatial and temporal scales
 - Decades: forest buffer restoration



1980





2015































Cyber infrastructure and tools










WikiWatershed

https://wikiwatershed.org/



Research at local and global scales






- 1. Stroud Water Research Center
- 2. Current research on PFAS
- 3. Questions and Discussion



What are PFAS?


Per and polyfluoroalkyl substances

- Class of >12,000 anthropogenic chemicals.
- Intentional and unintentional products of fluoropolymer synthesis.
 - Global production >230,000 tons/year
 - (Global pesticide production ~2.5M tons/year)
- Used in hundreds of applications.
 - Surfactant properties
 - High chemical and thermal stability



Source: Gulge et al. Environ. Sci.: Processes Impacts, 2020, 22, 2345–2373


PFAS toxicity to wildlife and humans



- Reproductive
- Endocrine
- Carcinogenic
- Neurotoxic
- Dyslipidemic
- Immunotoxic
- Bioaccumulate and biomagnify in wildlife

PFAS movement


from points of production and use to farms soil and water



Biosolids use in the US


and their potential as a source of PFAS

- Based on US EPA annual biosolids reports from roughly 2,250 larger facilities in 41 states. In 2023:
- Biosolids produced: ~4 million dry metric tons (dmt).
- Applied to agricultural land: 1.24 million dmt (~30%)




Source: https://www.epa.gov/biosolids/basic-information-about-sewage-sludge-and-biosolids


Biosolids use in Pennsylvania and their potential as a source of PFAS



Agricultural parcels in PA permitted to receive biosolids applications (includes active and inactive sites).


Source: PA Department of Environmental Protection






 Most soils treated with biosolids had measurable concentrations of at least one PFAS compound.

















PFAS distribution based on biosolid source




 Biosolids from private applicators had more detectable compounds and in generally higher concentrations



PFAS in water from wells of PA farms


- Five farms* receiving biosolids had PFAS in well water at levels above the drinking water maximum contaminant level (MCL = 4 ng/L)
- Different PFAS profile
 - Well water dominated by shorter-chain (9
 C or less) compounds.
- A direct connection between biosolid application and groundwater contamination cannot be confirmed at this time.




- Pennsylvania Farm
- Biosolids applied annually
- 2000-2022
- Most of the farm occupies the watershed of a 2nd order tributary of the Schuylkill River




- Pennsylvania Farm
- · Biosolids applied annually
- 2000-2022
- Most of the farm occupies the watershed of a 2nd order tributary of the Schuylkill River




- Precursors only present in biosolids
 - Likely break down



- Long-chain mostly in soils but not in water
 - Hydrophobic
 - Likely "sticking" to soils



- Short-chain mostly in soils but not in water
 - Hydrophilic
 - Likely mobilizing with water



New research question

- What are temporal dynamics of the PFAS in the soil and stream compartments?
 - Long-chain compounds will mobilize via overland flow during storm events
 - (sorbed to suspended sediments =particulate fraction).




New research question

- Weekly stream water and sediment sample collection
- Storm sample collection
- Water sample analysis for:
 - Suspended sediments
 - Dissolved organic carbon
 - Anions
 - o PFAS (USEPA 1633) (40 compounds)
 - Total
 - Dissolved (by centrifugation)
 - Particulate (difference between total and dissolved)
- Sediment sample analysis for:
 - Organic matter content
 - o PFAS (USEPA 1633) (40 compounds)

QA/QS for sampling equipment, field and lab blanks



Preliminary Results from first storm









Preliminary Results from first storm

- Contribution of particulate fraction to total PFAS concentration increased with chain-length.
- PFOSA (precursor) dominated the particulate fraction.




Preliminary Results from first storm

- Overall contribution from the particulate fraction was small.
- Total PFAS export was ~5x higher at stormflow conditions.



What have we learned so far?



- Biosolids used in Pennsylvania can be a significant source of PFAS contamination to farming soils, adjacent surface waters and <u>potentially</u> local drinking water supplies.
- Not all biosolids are created equal, not a widespread problem.

 O What is/are the source(s) of the biosolids and what kind of waste does the sourcing WWTP(s) receive?
- Different PFAS compounds behave differently in the agroecosystem.
- Stormflow conditions may represent hot moments for export of PFAS to downstream ecosystems.

Acknowledgments

Staff scientists: Stephanie
 Bernasconi, Mike Gentile,
 Kristen McCarthy, David
 Montgomery & Cooper Reilly

Funding Agencies

- Foundation for Food and Agricultural Research
- US Department of Agriculture

Farmers for access to their farms



Questions?

Thank you!

doviedo@stroudcenter.org



Stay in Touch

Subscribe to our monthly e-news for updates on freshwater research, environmental education, and watershed restoration.

Scan code or go to stroudcenter.org/subscribe

Follow Us









